

PAINLEVÉ IV: ROOTS AND ZEROS

P.L. ROBINSON

ABSTRACT. We consider the (real) fourth Painlevé equation in which both parameters vanish, analyzing the square-roots of its solutions and paying special attention to their zeros.

INTRODUCTION

In [2] we offered elementary proofs for fundamental properties of the unique triple-zero solution to the first Painlevé equation. In [3] we treated in a similar fashion all solutions to the second Painlevé equation whose graphs pass through the origin. Here we consider aspects of what is arguably the next case: the fourth Painlevé equation, which was discovered by Gambier. The general form of this equation is

$$\frac{d^2w}{dz^2} = \frac{1}{2w} \left(\frac{dw}{dz} \right)^2 + \frac{3}{2}w^3 + 4zw^2 + 2(z^2 - \alpha)w + \frac{\beta}{w}$$

where α and β are parameters. As is suggested by its form, this equation is properly *complex*: in fact, each of its solutions is meromorphic in the plane, with simple poles of residue ± 1 ; see [1] and references therein. Note the presence of the dependent variable in denominators: this separates the fourth Painlevé equation from the first and second; of course, it engenders some complications.

In line with the setting of our previous papers, we shall consider the fourth Painlevé equation in purely real terms; moreover, we shall only consider the case in which $\alpha = 0$ and $\beta = 0$. Accordingly, our version of the fourth Painlevé equation (P IV) is

$$\ddot{s} = \frac{1}{2s} \dot{s}^2 + \frac{3}{2}s^3 + 4ts^2 + 2t^2s$$

to be solved for real s as a function of real t ; our preference here for $\dot{s}$ over s' as notation for the derivative is largely on account of the otherwise awkward s'^2 or $(s')^2$ for its square. If the solution s is strictly positive then its (positive) square-root $\sqrt{s}$ satisfies a second-order equation that is simpler than P IV in having no first-derivative term and no awkward denominators; in the opposite direction, the squares of nowhere-zero solutions to this simpler equation satisfy P IV. Those circumstances in which solutions to P IV or the simpler equation have (isolated) zeros call for separate handling. All of these matters are discussed under the sections *Square-roots* (on the differential equations themselves) and *Isolated zeros* (on the special handling of zeros); in our final section on *Remarks* we address some related issues without proof.

SQUARE-ROOTS

We begin with some elementary observations regarding our version of the fourth Painlevé equation, which we restate for the record as

$$(P) \quad \ddot{s} = \frac{1}{2s} \dot{s}^2 + \frac{3}{2}s^3 + 4ts^2 + 2t^2s,$$

where the ratio on the right side is to be understood as a limit when necessary.

Observe that $\mathbf{P}$ may be reformulated in a number of ways. First we may clear the awkward denominator, thus:

$$2s\ddot{s} - \dot{s}^2 = 3s^4 + 8ts^3 + 4t^2s^2.$$

Further, we may factor the right side, thus:

$$2s\ddot{s} - \dot{s}^2 = s^2(3s + 2t)(s + 2t).$$

Observe also that reversing the sign of the dependent variable leads to the equation

$$(\overline{\mathbf{P}}) \quad \ddot{s} = \frac{1}{2s}\dot{s}^2 + \frac{3}{2}s^3 - 4ts^2 + 2t^2s;$$

of course, sign-reversal in $\overline{\mathbf{P}}$ leads to $\mathbf{P}$ likewise. Incidentally, passage between $\mathbf{P}$ and $\overline{\mathbf{P}}$ may also be effected by reversal of the independent variable. Of course, $\overline{\mathbf{P}}$ admits reformulations akin to those for $\mathbf{P}$ itself.

Let us agree to write $\mathbb{P}$ for the set comprising all solutions to the Painlevé equation $\mathbf{P}$; when extra clarity is called for, we may write $\mathbb{P}(I)$ for the set comprising all solutions to $\mathbf{P}$ on the open interval (more generally, open set) $I \subseteq \mathbb{R}$. Similarly, we write $\overline{\mathbb{P}}$ for the set of all solutions to $\overline{\mathbf{P}}$ (on some interval, which we may indicate for clarity). We observed above that multiplication by -1 yields a bijection

$$\mathbb{P} \rightarrow \overline{\mathbb{P}} : s \mapsto -s;$$

also, that reversal of the independent variable yields a bijection from $\mathbb{P}(I)$ to $\overline{\mathbb{P}}(-I)$.

Now, let $s \in \mathbb{P}$ be a *strictly positive* solution to $\mathbf{P}$ and write $\sigma := \sqrt{s} = s^{1/2}$ for its positive square-root. Certainly, σ is twice-differentiable. Further, from $s = \sigma^2$ it follows that

$$\dot{s} = 2\sigma\dot{\sigma}$$

so that

$$\dot{\sigma}^2 = \frac{\dot{s}^2}{4\sigma^2} = \frac{\dot{s}^2}{4s}$$

and

$$\ddot{s} = 2\dot{\sigma}^2 + 2\sigma\ddot{\sigma} = \frac{\dot{s}^2}{2s} + 2\sigma\ddot{\sigma}$$

so that

$$\ddot{s} - \frac{\dot{s}^2}{2s} = 2\sigma\ddot{\sigma}.$$

All of this requires only that the twice-differentiable function s be strictly positive. Recalling that s is a solution to $\mathbf{P}$ we deduce that

$$2\sigma\ddot{\sigma} = \ddot{s} - \frac{\dot{s}^2}{2s} = \frac{3}{2}s^3 + 4ts^2 + 2t^2s = \frac{1}{2}s(3s + 2t)(s + 2t)$$

or

$$\ddot{\sigma} = \frac{1}{4}\sigma(3\sigma^2 + 2t)(\sigma^2 + 2t).$$

This finding prompts us to formalize the auxiliary differential equation

$$(\mathbf{P}^{1/2}) \quad 4\ddot{\sigma} = \sigma(3\sigma^2 + 2t)(\sigma^2 + 2t)$$

alongside its companion

$$(\overline{\mathbf{P}}^{1/2}) \quad 4\ddot{\sigma} = \sigma(3\sigma^2 - 2t)(\sigma^2 - 2t).$$

It also prompts us to introduce $\mathbb{P}^{1/2}$ and $\overline{\mathbb{P}}^{1/2}$ for the corresponding spaces of solutions. Notice that the map $\sigma \mapsto -\sigma$ preserves the spaces $\mathbb{P}^{1/2}$ and $\overline{\mathbb{P}}^{1/2}$ while the map $t \mapsto -t$ interchanges them.

The following result was established in the motivating lead-up to equation $\mathbf{P}^{1/2}$.

Theorem 1. *If $s \in \mathbb{P}$ is strictly positive then $\sqrt{s} \in \mathbb{P}^{1/2}$.*

□

As a companion result, if $s \in \overline{\mathbb{P}}$ is strictly positive then a parallel argument places $\sqrt{s}$ in $\overline{\mathbb{P}}^{1/2}$; consequently, if $s \in \mathbb{P}$ is strictly *negative* then $\sqrt{-s} \in \overline{\mathbb{P}}^{1/2}$.

In the opposite direction, let $\sigma \in \mathbb{P}^{1/2}$ and write $s = \sigma^2$. Direct calculation as for the lead-up to $\mathbf{P}^{1/2}$ yields

$$2s\ddot{s} - \dot{s}^2 = 4\sigma^3\ddot{\sigma} = \sigma^4(3\sigma^2 + 2t)(\sigma^2 + 2t) = s^2(3s + 2t)(s + 2t).$$

If σ is never zero then we may divide by $2s$ throughout to see that $s \in \mathbb{P}$. We have established the following result.

Theorem 2. *If $\sigma \in \mathbb{P}^{1/2}$ is nowhere zero then $\sigma^2 \in \mathbb{P}$.*

□

Similarly, if $\sigma \in \overline{\mathbb{P}}^{1/2}$ is nowhere zero then $\sigma^2 \in \overline{\mathbb{P}}$ and $-\sigma^2 \in \mathbb{P}$ is strictly negative.

Throughout the present section, we have deliberately avoided situations in which $\sigma \in \overline{\mathbb{P}}^{1/2}$ or $s \in \mathbb{P}$ has a zero. We shall address such situations carefully in the next section; naturally, we may ignore the identically zero function.

ISOLATED ZEROS

As announced, we here consider situations in which $s \in \mathbb{P}$ or $\sigma \in \overline{\mathbb{P}}^{1/2}$ has a zero. Specifically, we shall assume that such a function has an isolated zero at the point a in the open interval I : more specifically, we shall assume that the function vanishes at a but at no other point of I . We wish to explore the extendibility of Theorem 1 and Theorem 2 to this context.

Observe at once from $\mathbf{P}$ (say in a reformulation) that if $s \in \mathbb{P}$ satisfies $s(a) = 0$ then automatically $\dot{s}(a) = 0$. In like but more straightforward manner, $\mathbf{P}^{1/2}$ tells us that if $\sigma \in \mathbb{P}^{1/2}$ satisfies $\sigma(a) = 0$ then automatically $\ddot{\sigma}(a) = 0$. We shall use these observations throughout the subsequent discussion, perhaps without comment.

Before proceeding further, it is convenient to draw attention to an important difference between $\mathbf{P}$ and $\mathbf{P}^{1/2}$. On the one hand, $\mathbf{P}^{1/2}$ has the form $\ddot{\sigma} = \Phi(t, \sigma)$ in which $\Phi(t, \sigma)$ is a polynomial; consequently, the initial value problem for $\mathbf{P}^{1/2}$ has a standard local existence-uniqueness theorem. On the other hand, $\mathbf{P}$ has the form $\ddot{s} = F(t, s, \dot{s})$ in which $F(t, s, \dot{s})$ is rational but has s in the denominator; the standard local existence-uniqueness theorem breaks down for initial data involving a zero of s . In fact, we have seen that if s satisfies $\mathbf{P}$ then the vanishing of $s(a)$ forces that of $\dot{s}(a)$; were standard local uniqueness to apply, a solution to $\mathbf{P}$ with a zero would vanish throughout its interval of definition.

Theorem 3. *Let $\sigma \in \mathbb{P}^{1/2}$ and let $\sigma \geq 0$ on $I \ni a$. If $\sigma(a) = 0$ then $\sigma = 0$.*

Proof. The hypotheses ensure that not only $\sigma(a) = 0$ but also $\dot{\sigma}(a) = 0$. The identically zero function satisfies $\mathbf{P}^{1/2}$ on I with the same initial data. The local uniqueness theorem for $\mathbf{P}^{1/2}$ now ensures that $\sigma = 0$. □

It follows at once that Theorem 1 has no direct extension allowing an isolated zero.

Theorem 4. *If $s \in \mathbb{P}$ is strictly positive except for an isolated zero at $a \in I$ then $\sqrt{s} \notin \mathbb{P}^{1/2}$.*

Proof. The (positive) square-root $\sqrt{s}$ is zero at $a \in I$ but strictly positive on $I \setminus \{a\}$; Theorem 3 therefore excludes $\sqrt{s}$ from $\mathbb{P}^{1/2}$. □

Notwithstanding this negative result, we have the following.

Theorem 5. *If $s \in \mathbb{P}$ is strictly positive except for an isolated zero at $a \in I$ then there exists $\sigma \in \mathbb{P}^{1/2}$ such that $s = \sigma^2$.*

Proof. To the left of a there are only two continuous square-roots of s , namely $\pm\sqrt{s}$; likewise to the right of a . Since the taking of like signs on each side of a leads to failure, we mix signs: thus, well-define σ on I by

$$\sigma(t) = \begin{cases} -\sqrt{s(t)} & \text{if } I \ni t \leq a, \\ +\sqrt{s(t)} & \text{if } I \ni t \geq a. \end{cases}$$

Theorem 1 easily places $\pm\sigma$ in $\mathbb{P}^{1/2}$ on $I \setminus \{a\}$; we must show that σ is twice-differentiable at its zero a with $\ddot{\sigma}(a) = 0$. Let $I \ni t \neq a$: as $s = \sigma^2$,

$$\dot{\sigma}(t) = \frac{\dot{s}(t)}{2\sigma(t)}$$

while as $s \in \mathbb{P}$ and $s(a) = 0$,

$$\frac{\dot{s}(t)^2}{2s(t)} = \ddot{s}(t) - \frac{1}{2}s(t)(3s(t) + 2t)(s(t) + 2t)$$

and

$$\lim_{t \rightarrow a} \frac{\dot{s}(t)^2}{4\sigma(t)^2} = \lim_{t \rightarrow a} \frac{\dot{s}(t)^2}{4s(t)} = \frac{1}{2}\ddot{s}(a).$$

Checking signs, the taking of square-roots yields

$$\lim_{t \rightarrow a} \dot{\sigma}(t) = \sqrt{\frac{1}{2}\ddot{s}(a)};$$

as σ is continuous on I , it follows that σ is (continuously) differentiable throughout I by an application of the mean value theorem. As σ satisfies $\mathbf{P}^{1/2}$ on $I \setminus \{a\}$, it follows that

$$\lim_{t \rightarrow a} \ddot{\sigma}(t) = \lim_{t \rightarrow a} \frac{1}{4}\sigma(t)(3\sigma(t)^2 + 2t)(\sigma(t)^2 + 2t) = 0$$

whence a further application of the mean value theorem to the continuous function $\dot{\sigma}$ shows that σ is twice-differentiable at a with $\ddot{\sigma}(a) = 0$ as required. $\square$

Of course, a similar argument shows that if $s \in \mathbb{P}$ is strictly negative except for an isolated zero at $a \in I$ then there exists $\sigma \in \overline{\mathbb{P}}^{1/2}$ such that $s = -\sigma^2$; again, σ takes opposite signs on opposite sides of a .

Theorem 4 and Theorem 5 are complements to Theorem 1 for cases in which $s \in \mathbb{P}$ has an isolated zero. There are analogous complements to Theorem 2 for cases in which $\sigma \in \mathbb{P}^{1/2}$ has an isolated zero.

The appropriate counterpart of Theorem 4 is immediate.

Theorem 6. *If $\sigma \in \mathbb{P}^{1/2}$ is strictly positive except for an isolated zero at $a \in I$ then $\sigma^2 \notin \mathbb{P}$.*

Proof. If σ^2 were to lie in $\mathbb{P}$ then its positive square-root would lie outside $\mathbb{P}^{1/2}$ according to Theorem 4; but this positive square-root is σ itself. $\square$

The appropriate counterpart of Theorem 5 requires just a little more work.

Theorem 7. *If $\sigma \in \mathbb{P}^{1/2}$ takes opposite signs on opposite sides of $a \in I$ then $\sigma^2 \in \mathbb{P}$.*

Proof. The twice-differentiable square $s := \sigma^2$ satisfies $\mathbf{P}$ on $I \setminus \{a\}$ by Theorem 2. Notice that if $t \in I$ then $\dot{s}(t) = 2\sigma(t)\dot{\sigma}(t)$ and

$$\ddot{s}(t) = 2\sigma(t)\ddot{\sigma}(t) + 2\dot{\sigma}(t)^2.$$

Consequently, as $\sigma(a) = 0$ it follows that

$$\ddot{s}(a) - \frac{1}{2}s(a)(3s(a) + 2a)(s(a) + 2a) = 2\dot{\sigma}(a)^2$$

and

$$\lim_{t \rightarrow a} \frac{\dot{s}(t)^2}{2s(t)} = \lim_{t \rightarrow a} \frac{(2\sigma(t)\dot{\sigma}(t))^2}{2\sigma(t)^2} = \lim_{t \rightarrow a} 2\dot{\sigma}(t)^2 = 2\dot{\sigma}(a)^2.$$

This shows that s satisfies **P** at a also and concludes the demonstration. $\square$

We close by remarking that the case of $s \in \mathbb{P}$ with an isolated zero at which the second derivative also vanishes is as tidy as can be: in fact, the case does not arise! Our first step towards this result is perhaps a little peculiar in hindsight.

Theorem 8. *Let $s \in \mathbb{P}$ have an isolated zero at a . If $\ddot{s}(a) = 0$ then each derivative of s vanishes at a .*

Proof. The second reformulation of **P** informs us that

$$2s\ddot{s} - \dot{s}^2 = s^2(3s + 2t)(s + 2t) = s^2Q$$

say, where Q is quadratic in s and t . Away from the isolated zero, we may differentiate: the resulting terms $\pm 2\dot{s}\ddot{s}$ on the left cancel, to yield

$$2s\ddot{\ddot{s}} = 2s\dot{s}Q + s^2\dot{Q}$$

whence

$$2\ddot{\ddot{s}} = 2\dot{s}Q + s\dot{Q}$$

away from a and hence at a also. All that remains is to differentiate inductively. $\square$

We can now see that this case is indeed vacuous: taking the (*difficult*!) meromorphicity of s for granted, the identity theorem implies that s is zero throughout the open interval in which a is an isolated zero; this is absurd!

In particular, it follows that $s \in \mathbb{P}$ cannot change sign at an isolated zero.

REMARKS

We round out our account with some miscellaneous comments on related topics of interest.

Recall that the fourth Painlevé equation is properly a complex differential equation. The process of passing to a square-root is naturally more elaborate in the complex setting: as we mentioned, solutions to the fourth Painlevé equation are meromorphic, with *simple* poles; square-roots of such functions cannot be meromorphic! Nonetheless, there is sufficient reason for further study of the relevant auxiliary equation

$$4\frac{d^2\omega}{dz^2} = \omega(3\omega^2 + 2z)(\omega^2 + 2z).$$

We have seen in our study of the fourth Painlevé equation **P** that the auxiliary differential equation **P**^{1/2} is of definite theoretical interest. In fact, this auxiliary equation is also of considerable practical help, aside from its ability to handle initial data involving a zero. We began exploring solutions of the fourth Painlevé equation with the aid of WZGrapher, a valuable free-ware program developed by Walter Zorn. Quite early in our explorations, we noticed apparent graphical instabilities: for example, solutions of **P** with certain initial data would at first appear to be oscillatory; upon zooming out, such a solution might seem to suffer a catastrophe, oscillations disappearing and being replaced by a blow-up or spike; upon zooming out further, oscillations might reappear; and so on. Not surprisingly, such catastrophic behaviour manifests itself at a zero of the solution and so involves the awkward denominator in **P**. These apparent graphical instabilities seem to be removed by passage to the corresponding solutions of **P**^{1/2}, as the reader may care to see using WZGrapher.

The factorized form of the fourth Painlevé equation $\mathbf{P}$:

$$\ddot{s} - \frac{\dot{s}^2}{2s} = \frac{1}{2}s(3s + 2t)(s + 2t)$$

indicates that the lines ' $s = 0$ ', ' $s = -2t/3$ ' and ' $s = -2t$ ' have geometric significance for its solutions. Similarly, ' $\sigma = 0$ ' and the parabolas ' $\sigma^2 = -2t/3$ ' and ' $\sigma^2 = -2t$ ' have geometric significance for solutions to the auxiliary equation $\mathbf{P}^{1/2}$:

$$4\ddot{\sigma} = \sigma(3\sigma^2 + 2t)(\sigma^2 + 2t).$$

This geometric significance can be seen in a concavity diagram. The curves ' $\sigma = 0$ ', ' $\sigma^2 = -2t/3$ ' and ' $\sigma^2 = -2t$ ' divide the (t, σ) -plane into regions. The sign of $\ddot{\sigma}$ is negative/positive in the regions directly above/below the half-parabola ' $\sigma = +\sqrt{-2t/3}$ ' so that solutions to $\mathbf{P}^{1/2}$ have the opportunity to oscillate about this half-parabola; similarly, solutions to $\mathbf{P}^{1/2}$ may oscillate about ' $\sigma = -\sqrt{-2t/3}$ '.

In fact, experimentation with WZGrapher reveals that solutions to $\mathbf{P}^{1/2}$ that do not suffer blow-up in both time directions tend to display steadily decaying oscillations about the upper or lower half of the parabola ' $\sigma^2 = -2t/3$ ' as $t \rightarrow -\infty$; and that solutions often tend to linger alongside ' $\sigma = 0$ ' or ' $\sigma^2 = -2t$ ' as they make more or less extended approaches to tangency. Also, it not infrequently happens that a minuscule change in initial data causes a solution σ to flip its oscillations from one half-parabola to the other, or to flip the direction of its finite-time blow-up, in such a way that the sudden transition is not detectable in σ^2 . On a more aesthetic note, when oscillations of $\sigma \in \mathbb{P}^{1/2}$ occur about a half-parabola ' $\sigma = \pm\sqrt{-2t/3}$ ' they are quite evenly balanced. By contrast, when oscillations of $s \in \mathbb{P}$ occur about the line ' $s = -2t/3$ ' they are uneven, displaying larger arches on the side of the line away from ' $s = 0$ '. Of course, squaring accounts for the difference.

One relatively simple family of illustrative examples takes $\sigma \in \mathbb{P}^{1/2}$ with $\sigma(0) = 0$ and $\dot{\sigma}(0)$ strictly positive. As $\dot{\sigma}(0)$ increases from 0 to a little beyond 1.169868591, two gradual changes to the solution σ take place simultaneously: on the one hand, σ oscillates about ' $\sigma = -\sqrt{-2t/3}$ ', the amplitude of the oscillations initially decreasing and finally increasing; on the other hand, σ lingers initially along ' $\sigma = 0$ ' and finally along ' $\sigma = -\sqrt{-2t}$ '; when $\dot{\sigma}(0)$ is around 0.65 the oscillations have their least amplitude and there is no lingering along either curve. As $\dot{\sigma}(0)$ increases from 1.169868591 to 1.169868592 the oscillations disappear, to be replaced by a negative blow-up in finite negative time; thereafter, the lingering along ' $\sigma = -\sqrt{-2t}$ ' gradually disappears and the finite-time blow-up accelerates. Throughout, $\sigma \in \mathbb{P}^{1/2}$ has a unique zero, at which it changes sign; accordingly, its square lies in $\mathbb{P}$.

We leave to the reader the pleasure of exploring this family of examples in WZGrapher (or some similar program). Among many other families to explore, we recommend the following: take $\sigma(0) = 1$ and let $\dot{\sigma}(0)$ run from -0.933899363 to 1.579186627 , noting the several transitions with reference to ' $\sigma = 0$ ', ' $\sigma^2 = -2t/3$ ' and ' $\sigma^2 = -2t$ '; take $\sigma(-6) = 2$ and let $\dot{\sigma}(-6)$ run from -0.170889967 to -0.170889968 (!).

REFERENCES

- [1] V.I. Gromak, I. Laine and S. Shimomura, *Painlevé Differential Equations in the Complex Plane*, de Gruyter (2002).
- [2] P.L. Robinson, *The Triple-Zero Painlevé I Transcendent*, arXiv 1607.07088 (2016).
- [3] P.L. Robinson, *Homogeneous Painlevé II Transcendents*, arXiv 1608.02139 (2016).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, GAINESVILLE FL 32611 USA
E-mail address: paulr@ufl.edu